

Small index subgroups of the mapping class group

LUIS PARIS

February 2, 2008

Abstract

We prove that the mapping class group of a closed oriented surface of genus $\rho \geq 3$ has no proper subgroup of index $\leq 4\rho + 4$.

AMS Subject Classification. Primary: 57M99. Secondary: 20F36.

Let Σ be a closed oriented surface of genus $\rho \geq 1$, and let $\text{Homeo}^+(\Sigma)$ be the group of homeomorphisms $h : \Sigma \rightarrow \Sigma$ that preserve the orientation. The *mapping class group* of Σ is defined to be the group $\mathcal{M}(\Sigma) = \pi_0(\text{Homeo}^+(\Sigma))$ of isotopy classes of elements of $\text{Homeo}^+(\Sigma)$. This group plays an important role in low-dimensional topology, in Teichmüller theory, and in group theory. We refer to [6] for a general survey on the subject, and turn to the main object of the present paper: the finite index subgroups of $\mathcal{M}(\Sigma)$.

A group G is called *residually finite* if, for all $g \in G \setminus \{1\}$, there exists a homomorphism $\varphi : G \rightarrow H$ such that H is finite and $\varphi(g) \neq 1$. By [4], the mapping class group $\mathcal{M}(\Sigma)$ is residually finite. This means that $\mathcal{M}(\Sigma)$ has many finite index subgroups, numerous enough to separate its elements. Nevertheless, we are going to prove:

Theorem 1. *Let Σ be a closed oriented surface of genus $\rho \geq 3$. Then $\mathcal{M}(\Sigma)$ has no proper subgroup of index $\leq 4\rho + 4$.*

Remark. If $\rho = 1$, then $H_1(\mathcal{M}(\Sigma)) \simeq \mathbb{Z}/12\mathbb{Z}$, thus $\mathcal{M}(\Sigma)$ has (normal) subgroups of index 2 and 3. If $\rho = 2$, then $H_1(\mathcal{M}(\Sigma)) \simeq \mathbb{Z}/10\mathbb{Z}$ (see [9]), thus $\mathcal{M}(\Sigma)$ has (normal) subgroups of index 2 and 5. So, the hypothesis $\rho \geq 3$ is needed in Theorem 1. Note also that the statement is valid for all the subgroups, not only for the normal ones.

The rest of the article is dedicated to the proof of Theorem 1. So, from now on, Σ denotes a closed oriented surface of genus $\rho \geq 3$.

The two main tools that we use in our proof are the braid groups and the Dehn twists. For $n \geq 2$, we denote by $\mathcal{B}_n$ the braid group on n strands. This has a presentation with generators $\sigma_1, \dots, \sigma_{n-1}$ and relations

$$\begin{aligned} \sigma_i \sigma_j &= \sigma_j \sigma_i & \text{if } |i - j| \geq 2, \\ \sigma_i \sigma_j \sigma_i &= \sigma_j \sigma_i \sigma_j & \text{if } |i - j| = 1. \end{aligned}$$

The Dehn twist along an essential circle $a : \mathbb{S}^1 \hookrightarrow \Sigma$ will be denoted by τ_a . We assume the reader to be familiar with these two concepts, and we refer to [2] for a detailed text on them.

The following is easy to prove and can be found, for instance, in [3].

Lemma 2. *Let a, b be two essential circles which intersect transversely. Then*

$$\begin{aligned} \tau_a \tau_b &= \tau_b \tau_a & \text{if } a \cap b = \emptyset, \\ \tau_a \tau_b \tau_a &= \tau_b \tau_a \tau_b & \text{if } |a \cap b| = 1. \end{aligned}$$

Now, consider the essential circles $a_0, a_1, \dots, a_{2\rho+1}$ drawn in Figure 1, and, for $0 \leq i \leq 2\rho + 1$, let τ_i denote the Dehn twist along a_i .




Figure 1. Essential circles in Σ .

Proposition 3.

1. *The mapping $\sigma_i \mapsto \tau_i$, $1 \leq i \leq 2\rho + 1$, induces a homomorphism $\iota : \mathcal{B}_{2\rho+2} \rightarrow \mathcal{M}(\Sigma)$. Furthermore, we have the relations*

$$\begin{aligned} \tau_0 \tau_i &= \tau_i \tau_0 & \text{for } 1 \leq i \leq 2\rho + 1 \text{ and } i \neq 4, \\ \tau_0 \tau_4 \tau_0 &= \tau_4 \tau_0 \tau_4. \end{aligned}$$

2. (Humphries [5]) *The Dehn twists $\tau_0, \tau_1, \dots, \tau_{2\rho}$ generate $\mathcal{M}(\Sigma)$.*
3. (Powell [10]) *$H_1(\mathcal{M}(\Sigma)) = \{0\}$.*

Remark. Proposition 3.1 is a straightforward consequence of Lemma 2. Note also that $\iota : \mathcal{B}_{2\rho+2} \rightarrow \mathcal{M}(\Sigma)$ is not injective (see [7]).

For $k \geq 1$, we denote by Sym_k the symmetric group of $\{1, 2, \dots, k\}$. Let G be a group. A homomorphism $\varphi : G \rightarrow \text{Sym}_k$ is called *transitive* if $\text{Im } \varphi$ acts transitively on $\{1, 2, \dots, k\}$. If $\varphi : G \rightarrow \text{Sym}_k$ is a transitive homomorphism, then $\text{Stab}_\varphi(1) = \{g \in G; \varphi(g)(1) = 1\}$ is an index k subgroup of G . Reciprocally, if H is an index k subgroup of G , then there exists a transitive homomorphism $\varphi : G \rightarrow \text{Sym}_k$ such that $H = \text{Stab}_\varphi(1)$ (take the action of G on the left cosets of H). So, in order to prove Theorem 1, it suffices to prove that there is no transitive homomorphism $\varphi : \mathcal{M}(\Sigma) \rightarrow \text{Sym}_k$ for any $2 \leq k \leq 2\rho + 4$.

For $u \in \text{Sym}_k$, we denote by $\text{Ad}_u : \text{Sym}_k \rightarrow \text{Sym}_k$ the inner automorphism which sends v to uvu^{-1} for all $v \in \text{Sym}_k$. We say that two homomorphisms $\varphi, \varphi' : G \rightarrow \text{Sym}_k$ are *conjugate* if there exists $u \in \text{Sym}_k$ such that $\varphi' = \text{Ad}_u \circ \varphi$. To be conjugate is an equivalence relation that we denote by $\varphi \sim \varphi'$. The *product* of two homomorphisms $\varphi_1 : G \rightarrow \text{Sym}_{k_1}$ and $\varphi_2 : G \rightarrow \text{Sym}_{k_2}$ is the homomorphism $\varphi_1 \times \varphi_2 : G \rightarrow \text{Sym}_{k_1+k_2}$ defined by

$$(\varphi_1 \times \varphi_2)(g)(i) = \begin{cases} \varphi_1(g)(i) & \text{if } 1 \leq i \leq k_1, \\ \varphi_2(g)(i - k_1) + k_1 & \text{if } k_1 + 1 \leq i \leq k_1 + k_2. \end{cases}$$

Clearly, for every homomorphism $\varphi : G \rightarrow \text{Sym}_k$, there exists a partition $(k_1, k_2, \dots, k_p)$ of k and a collection $\{\varphi_j : G \rightarrow \text{Sym}_{k_j}; 1 \leq j \leq p\}$ of transitive homomorphisms such that

$$\varphi \sim \varphi_1 \times \varphi_2 \times \dots \times \varphi_p.$$

Moreover, this decomposition is unique up to a permutation of the factors.

Using the above defined product of homomorphisms, it is easily shown that the existence of a transitive homomorphism $\varphi : \mathcal{M}(\Sigma) \rightarrow \text{Sym}_k$ for some $2 \leq k \leq 4\rho + 4$ is equivalent to the existence of a non-trivial homomorphism $\varphi : \mathcal{M}(\Sigma) \rightarrow \text{Sym}_{4\rho+4}$.

We set $n = 2\rho+2$ and we turn now to give the classification of the homomorphisms from $\mathcal{B}_n$ to Sym_{2n} (see Corollary 7). Then, given a homomorphism $\varphi : \mathcal{M}(\Sigma) \rightarrow \text{Sym}_{2n} = \text{Sym}_{4\rho+4}$, our strategy for proving that φ is trivial will consist on exploring the different possibilities for the composition $\psi = \varphi \circ \iota : \mathcal{B}_n \rightarrow \text{Sym}_{2n}$, where $\iota : \mathcal{B}_n = \mathcal{B}_{2\rho+2} \rightarrow \mathcal{M}(\Sigma)$ is the homomorphism of Proposition 3.1.

Let $k \in \mathbb{N}$ and let $w \in \text{Sym}_k$. Then the mapping $\sigma_i \mapsto w$, $1 \leq i \leq k$, induces a homomorphism $\text{cycl}_w : \mathcal{B}_k \rightarrow \text{Sym}_k$ called a *cyclic homomorphism*. Note that cycl_w is transitive if and only if w is a cycle of length k . On the other hand, the mapping $\sigma_i \mapsto (i, i+1) \in \text{Sym}_k$, $1 \leq i \leq k-1$, induces a transitive homomorphism $\rho_S : \mathcal{B}_k \rightarrow \text{Sym}_k$ called the *standard homomorphism*.

Theorem 4 (Artin [1]). *Let $n \geq 7$.*

1. For $k < n$, every transitive homomorphism $\psi : \mathcal{B}_n \rightarrow \text{Sym}_k$ is cyclic.
2. Every transitive homomorphism $\psi : \mathcal{B}_n \rightarrow \text{Sym}_n$ is either cyclic or conjugate to the standard homomorphism ρ_S .

Remark. For $n = 4$ or 6 , there are transitive homomorphisms $\mathcal{B}_n \rightarrow \text{Sym}_n$ which are neither cyclic, nor conjugate to ρ_S : there are 3 for $n = 4$, and 1 for $n = 6$, up to conjugation (see [1]). However, in this paper we are concerned only by the case $n \geq 8$.

Proposition 5 (Lin [8]). *Let $n \geq 7$.*

1. *The mapping*

$$\sigma_i \mapsto (i, n+i+1, n+i, i+1), \quad 1 \leq i \leq n-1,$$

induces a transitive homomorphism $\rho_{L1} : \mathcal{B}_n \rightarrow \text{Sym}_{2n}$.

2. *The mapping*

$$\sigma_i \mapsto \prod_{j=1}^{i-1} (j, n+j) \cdot (i, n+i+1, n+i, i+1) \cdot \prod_{j=i+2}^n (j, n+j), \quad 1 \leq i \leq n-1,$$

induces a transitive homomorphism $\rho_{L2} : \mathcal{B}_n \rightarrow \text{Sym}_{2n}$.

3. *The mapping*

$$\sigma_i \mapsto \prod_{j=1}^{i-1} (j, n+j) \cdot (i, n+i+1)(i+1, n+i) \cdot \prod_{j=i+2}^n (j, n+j), \quad 1 \leq i \leq n-1,$$

induces a transitive homomorphism $\rho_{L3} : \mathcal{B}_n \rightarrow \text{Sym}_{2n}$.

The homomorphisms $\rho_{L1}, \rho_{L2}, \rho_{L3}$ of Proposition 5 will be called *Lin homomorphisms*.

Theorem 6 (Lin [8]). *Let $n \geq 7$.*

1. *For $n+1 \leq k \leq 2n-1$, every transitive homomorphism $\psi : \mathcal{B}_n \rightarrow \text{Sym}_k$ is cyclic.*
2. *Every transitive homomorphism $\psi : \mathcal{B}_n \rightarrow \text{Sym}_{2n}$ is either cyclic or conjugate to some ρ_{Lj} , where $j \in \{1, 2, 3\}$.*

Corollary 7. *Let $n \geq 7$. Let $\psi : \mathcal{B}_n \rightarrow \text{Sym}_{2n}$ be a homomorphism. Then either*

- (a) *ψ is cyclic; or*
- (b) *$\psi \sim \rho_S \times \text{cycl}_w$ for some $w \in \text{Sym}_n$; or*

(c) $\psi \sim \rho_S \times \rho_S$; or

(d) $\psi \sim \rho_{Lj}$ for some $j \in \{1, 2, 3\}$.

Proof of Theorem 1. Set $n = 2\rho + 2 \geq 8$. Let $\varphi : \mathcal{M}(\Sigma) \rightarrow \text{Sym}_{4\rho+4} = \text{Sym}_{2n}$ be a homomorphism. As pointed out before, we must prove that φ is trivial. We set $\psi = \varphi \circ \iota : \mathcal{B}_n \rightarrow \text{Sym}_{2n}$, where $\iota : \mathcal{B}_n \rightarrow \mathcal{M}(\Sigma)$ is the homomorphism of Proposition 3.1. By Corollary 7, we have either

(a) $\psi = \text{cycl}_w$ for some $w \in \text{Sym}_{2n}$; or

(b) $\psi \sim \rho_S \times \text{cycl}_w$ for some $w \in \text{Sym}_n$; or

(c) $\psi \sim \rho_S \times \rho_S$; or

(d) $\psi \sim \rho_{Lj}$ for some $j \in \{1, 2, 3\}$.

Case 1: Suppose $\psi = \text{cycl}_w$, where $w \in \text{Sym}_{2n}$.

Set $f = \tau_4\tau_0$ and observe that $f\tau_4f^{-1} = \tau_0$, and $f\tau_1f^{-1} = \tau_1$. We have $\varphi(\tau_i) = \psi(\sigma_i) = w$ for all $1 \leq i \leq 2\rho$. Moreover,

$$\varphi(\tau_0) = \varphi(f)\varphi(\tau_4)\varphi(f^{-1}) = \varphi(f)\varphi(\tau_1)\varphi(f^{-1}) = \varphi(\tau_1) = w.$$

Since $\mathcal{M}(\Sigma)$ is generated by $\tau_0, \tau_1, \dots, \tau_{2\rho}$ (see Proposition 3.2), it follows that the image of φ is the cyclic group generated by w : an abelian group. Since $H_1(\mathcal{M}(\Sigma)) = \{0\}$ (see Proposition 3.3), we conclude that $\text{Im } \varphi$ is trivial, that is, $w = 1$ and φ is trivial.

Case 2: Suppose $\psi = \rho_S \times \text{cycl}_w$, where $w \in \text{Sym}_n$.

For $v \in \text{Sym}_{2n}$, we define the *support* of v to be $\text{Supp}(v) = \{i \in \{1, \dots, 2n\}; v(i) \neq i\}$. We will often use the fact that two elements commute if their supports are disjoint.

Let $w_0 \in \text{Sym}_{2n}$ be defined by

$$w_0(i) = \begin{cases} i & \text{if } 1 \leq i \leq n, \\ w(i-n) + n & \text{if } n+1 \leq i \leq 2n. \end{cases}$$

So, $\text{Supp}(w_0) \subset \{n+1, n+2, \dots, 2n\}$, and

$$\varphi(\tau_i) = \psi(\sigma_i) = (i, i+1)w_0, \quad \text{for all } 1 \leq i \leq n-1.$$

Set $u_0 = \varphi(\tau_0)$. The permutation u_0 commutes with $\varphi(\tau_1)\varphi(\tau_2)^{-1} = (1, 2, 3)$, with $\varphi(\tau_2)\varphi(\tau_3)^{-1} = (2, 3, 4)$, and with $\varphi(\tau_1)\varphi(\tau_3)^{-1} = (1, 2)(3, 4)$, thus

$$u_0(\{1, 2, 3\}) = \{1, 2, 3\}, \quad u_0(\{2, 3, 4\}) = \{2, 3, 4\},$$

and either

$$\begin{aligned} u_0(\{1, 2\}) = \{1, 2\} \quad \text{and} \quad u_0(\{3, 4\}) = \{3, 4\}, \quad \text{or} \\ u_0(\{1, 2\}) = \{3, 4\} \quad \text{and} \quad u_0(\{3, 4\}) = \{1, 2\}. \end{aligned}$$

This clearly implies that $u_0(i) = i$ for all $1 \leq i \leq 4$. Similarly, the fact that u_0 commutes with $\varphi(\tau_i) = (i, i+1)w_0$ for all $5 \leq i \leq n-1$ implies that $u_0(i) = i$ for all $5 \leq i \leq n$. So, $\text{Supp}(u_0) \subset \{n+1, \dots, 2n\}$. Now, from the equality $\tau_0\tau_4\tau_0 = \tau_4\tau_0\tau_4$ follows

$$(4, 5)u_0w_0u_0 = (4, 5)(4, 5)w_0u_0w_0 = w_0u_0w_0,$$

and this last equality cannot hold: a contradiction.

Case 3: Suppose $\psi = \rho_S \times \rho_S$.

We have

$$\varphi(\tau_i) = \psi(\sigma_i) = (i, i+1)(n+i, n+i+1), \quad \text{for all } 1 \leq i \leq n-1.$$

Set $u_0 = \varphi(\tau_0)$. Recall that $\tau_0 = f\tau_4f^{-1}$, where $f = \tau_4\tau_0$, thus u_0 is conjugate to $\varphi(\tau_4) = (4, 5)(n+4, n+5)$, therefore u_0 is the disjoint product of two transpositions. Since u_0 commutes with $\varphi(\tau_i) = (i, i+1)(n+i, n+i+1)$ for all $1 \leq i \leq 3$, we have

$$u_0(\{i, i+1, n+i, n+i+1\}) = \{i, i+1, n+i, n+i+1\}$$

for all $1 \leq i \leq 3$, thus $u_0(\{i, n+i\}) = \{i, n+i\}$ for all $1 \leq i \leq 4$. If $u_0(i) = n+i$ then $u_0(i+1) = n+i+1$, since u_0 commutes with $\varphi(\tau_i) = (i, i+1)(n+i, n+i+1)$. Similarly, if $u_0(i) = i$, then $u_0(i+1) = i+1$. So, either $u_0(i) = i$ and $u_0(n+i) = n+i$ for all $1 \leq i \leq 4$, or $u_0(i) = n+i$ and $u_0(n+i) = i$ for all $1 \leq i \leq 4$. The latter case cannot hold because u_0 is the disjoint product of two transpositions, thus $u_0(i) = i$ and $u_0(n+i) = n+i$ for all $1 \leq i \leq 4$. But, the fact that u_0 commutes with $\varphi(\tau_i) = (i, i+1)(n+i, n+i+1)$ for all $5 \leq i \leq n-1$ also implies that $u_0(i) = i$ and $u_0(n+i) = n+i$ for all $5 \leq i \leq n$. So, $u_0 = \text{Id}$: a contradiction.

Case 4: Suppose $\psi = \rho_{L1}$.

Let $\text{sg} : \text{Sym}_{2n} \rightarrow \{\pm 1\}$ denote the signature. Observe that

$$(\text{sg} \circ \varphi)(\tau_1) = (\text{sg} \circ \psi)(\sigma_1) = \text{sg}((1, n+2, n+1, 2)) = -1,$$

thus $\text{sg} \circ \varphi : \mathcal{M}(\Sigma) \rightarrow \{\pm 1\}$ is surjective. This contradicts the fact that $H_1(\mathcal{M}(\Sigma)) = \{0\}$ (see Proposition 3.3).

Case 5: Suppose $\psi = \rho_{L2}$.

Note that $n = 2\rho + 2$ is even, thus

$$(\text{sg} \circ \varphi)(\tau_1) = (\text{sg} \circ \psi)(\sigma_1) = \text{sg} \left((1, n+2, n+1, 2) \cdot \prod_{j=3}^n (j, n+j) \right) = (-1)^{n-1} = -1,$$

therefore $\text{sg} \circ \varphi : \mathcal{M}(\Sigma) \rightarrow \{\pm 1\}$ is surjective. As pointed out in the previous case, such a homomorphism cannot exist because $H_1(\mathcal{M}(\Sigma)) = \{0\}$.

Case 6: Suppose $\psi = \rho_L 3$.

We have

$$\varphi(\tau_i) = \psi(\sigma_i) = \prod_{j=1}^{i-1} (j, n+j) \cdot (i, n+i+1)(i+1, n+i) \cdot \prod_{j=i+2}^n (j, n+j),$$

for all $1 \leq i \leq n-1$. Set $u_0 = \varphi(\tau_0)$. Recall that τ_0 is conjugate to τ_4 , thus u_0 is conjugate to $\varphi(\tau_4)$, therefore u_0 is the disjoint product of n transpositions. The permutation u_0 commutes with the following three:

$$\begin{aligned} \varphi(\tau_1)\varphi(\tau_2) &= (1, 2, 3)(n+1, n+2, n+3), \\ \varphi(\tau_1)\varphi(\tau_3) &= (1, 2)(3, 4)(n+1, n+2)(n+3, n+4), \\ \varphi(\tau_2)\varphi(\tau_3) &= (2, 3, 4)(n+2, n+3, n+4), \end{aligned}$$

thus

$$\begin{aligned} u_0(\{1, 2, 3, n+1, n+2, n+3\}) &= \{1, 2, 3, n+1, n+2, n+3\}, \\ u_0(\{1, 2, 3, 4, n+1, n+2, n+3, n+4\}) &= \{1, 2, 3, 4, n+1, n+2, n+3, n+4\}, \\ u_0(\{2, 3, 4, n+2, n+3, n+4\}) &= \{2, 3, 4, n+2, n+3, n+4\}, \end{aligned}$$

hence

$$u_0(\{1, n+1\}) = \{1, n+1\} \quad \text{and} \quad u_0(\{4, n+4\}) = \{4, n+4\}.$$

Moreover, the above two equalities plus the fact that u_0 commutes with $\varphi(\tau_1)\varphi(\tau_3) = (1, 2)(3, 4)(n+1, n+2)(n+3, n+4)$ imply that we also have

$$u_0(\{2, n+2\}) = \{2, n+2\} \quad \text{and} \quad u_0(\{3, n+3\}) = \{3, n+3\}.$$

Similarly, the fact that u_0 commutes with $\varphi(\tau_i)$ for all $5 \leq i \leq n-1$ implies that $u_0(\{i, n+i\}) = \{i, n+i\}$ for all $5 \leq i \leq n$. Finally, since u_0 is the disjoint product of n transpositions, we conclude that

$$u_0 = \prod_{j=1}^n (j, n+j).$$

Now, a direct calculation shows that

$$u_0\varphi(\tau_4)u_0 = \varphi(\tau_4) \neq u_0 = \varphi(\tau_4)u_0\varphi(\tau_4),$$

and this contradicts the equality $\tau_0\tau_4\tau_0 = \tau_4\tau_0\tau_4$ of Proposition 3.1. $\square$

References

- [1] **E. Artin.** *Braids and permutations.* Ann. of Math. (2) **48** (1947), 643–649.
- [2] **J.S. Birman.** *Braids, links, and mapping class groups.* Annals of Mathematics Studies, No. 82. Princeton University Press, Princeton, N.J., 1974.
- [3] **J.S. Birman.** *Mapping class groups of surfaces.* Braids (Santa Cruz, CA, 1986), 13–43, Contemp. Math., 78, Amer. Math. Soc., Providence, RI, 1988.
- [4] **E.K. Grossman.** *On the residual finiteness of certain mapping class groups.* J. London Math. Soc. (2) **9** (1974/75), 160–164.
- [5] **S.P. Humphries.** *Generators for the mapping class group.* Topology of low-dimensional manifolds (Proc. Second Sussex Conf., Chelwood Gate, 1977), pp. 44–47, Lecture Notes in Math., 722, Springer, Berlin, 1979.
- [6] **N.V. Ivanov.** *Mapping class groups.* Handbook of geometric topology, 523–633, North-Holland, Amsterdam, 2002.
- [7] **C. Labruère, L. Paris.** *Presentations for the punctured mapping class groups in terms of Artin groups.* Algebr. Geom. Topol. **1** (2001), 73–114.
- [8] **V. Lin.** *Braids and permutations.* Preprint, 2004. ArXiv: [math.GR/0404528](https://arxiv.org/abs/math/0404528).
- [9] **D. Mumford.** *Abelian quotients of the Teichmüller modular group.* J. Analyse Math. **18** (1967), 227–244.
- [10] **J. Powell.** *Two theorems on the mapping class group of a surface.* Proc. Amer. Math. Soc. **68** (1978), no. 3, 347–350.

Luis Paris,

Institut de Mathématiques de Bourgogne, UMR 5584 du CNRS, Université de Bourgogne,
B.P. 47870, 21078 Dijon cedex, France

E-mail: lparis@u-bourgogne.fr